

Mahler Discrete Residues and Summability for Rational Functions

Yi Zhang

Department of Foundational Mathematics
Xi'an Jiaotong-Liverpool University, Suzhou, China

Joint work with Carlos E. Arreche



Linear Mahler equations

Let \mathbb{K} be an algebraically closed field of char 0, x be an indeterminate, and $p \in \mathbb{Z}_{\geq 2}$.

Consider

$$\ell_r(x)y(x^{p^r}) + \ell_{r-1}(x)y(x^{p^{r-1}}) + \cdots + \ell_0(x)y(x) = f(x), \quad (1)$$

where $\ell_i, f \in \mathbb{K}[x]$ are given, $y(x)$ is unknown. A solution of (1) is called a **Mahler function**.

(Mahler 1929): study Mahler equations to prove the transcendence of values of some functions.

Fact: the generating series of any p -automatic sequence (such as the Baum–Sweet and the Rudin–Shapiro sequences) is a Mahler function.

Differential Galois Theory

Using differential Galois theory, we can determine the differential-algebraic relations between solutions of Mahler equations.

Example (Roques 2018): A Galoisian proof that the generating series of the Baum-Sweet and Rudin-Shapiro sequences are algebraic independent over $\bar{\mathbb{Q}}(x)$.

Goal: Design effective algorithms for computing differential Galois groups of a given linear Mahler equations.

Discrete residues, telescopers, and Galois theory

Endow $\mathbb{K}(x)$ with one of the $\sigma\delta$ -field structures:

(S) $\sigma : f(x) \mapsto f(x+1)$ and $\delta = \frac{d}{dx}$;

(Q) $\sigma : f(x) \mapsto f(qx)$ with $q \in \mathbb{K}^\times$ not root of unity and $\delta = x \frac{d}{dx}$.

Discrete residues, telescopers, and Galois theory

Endow $\mathbb{K}(x)$ with one of the $\sigma\delta$ -field structures:

(S) $\sigma : f(x) \mapsto f(x+1)$ and $\delta = \frac{d}{dx}$;

(Q) $\sigma : f(x) \mapsto f(qx)$ with $q \in \mathbb{K}^\times$ not root of unity and $\delta = x \frac{d}{dx}$.

Let $z_1, \dots, z_n \in F$, a $\sigma\delta$ -extension of $\mathbb{K}(x)$ with $F^\sigma = \mathbb{K}$, satisfying

$$\sigma(z_i) = a_i z_i \quad \text{for some } a_1, \dots, a_n \in \mathbb{K}(x)^\times.$$

Discrete residues, telescopers, and Galois theory

Endow $\mathbb{K}(x)$ with one of the $\sigma\delta$ -field structures:

(S) $\sigma : f(x) \mapsto f(x+1)$ and $\delta = \frac{d}{dx}$;

(Q) $\sigma : f(x) \mapsto f(qx)$ with $q \in \mathbb{K}^\times$ not root of unity and $\delta = x \frac{d}{dx}$.

Let $z_1, \dots, z_n \in F$, a $\sigma\delta$ -extension of $\mathbb{K}(x)$ with $F^\sigma = \mathbb{K}$, satisfying

$$\sigma(z_i) = a_i z_i \quad \text{for some } a_1, \dots, a_n \in \mathbb{K}(x)^\times.$$

Proposition (Hardouin-Singer 2008) z_1, \dots, z_n are δ -dependent over $\mathbb{K}(x)$ iff $\exists \mathcal{L}_1, \dots, \mathcal{L}_n \in \mathbb{K}[\delta]$, linear δ -operators with coefficients in \mathbb{K} , not all 0, and $g \in \mathbb{K}(x)$:

$$\mathcal{L}_1 \left(\frac{\delta(a_1)}{a_1} \right) + \dots + \mathcal{L}_n \left(\frac{\delta(a_n)}{a_n} \right) = \sigma(g) - g.$$

Discrete residues, telescopers, and Galois theory

Endow $\mathbb{K}(x)$ with one of the $\sigma\delta$ -field structures:

(S) $\sigma : f(x) \mapsto f(x+1)$ and $\delta = \frac{d}{dx}$;

(Q) $\sigma : f(x) \mapsto f(qx)$ with $q \in \mathbb{K}^\times$ not root of unity and $\delta = x \frac{d}{dx}$.

Let $z_1, \dots, z_n \in F$, a $\sigma\delta$ -extension of $\mathbb{K}(x)$ with $F^\sigma = \mathbb{K}$, satisfying

$$\sigma(z_i) = a_i z_i \quad \text{for some } a_1, \dots, a_n \in \mathbb{K}(x)^\times.$$

Proposition (Hardouin-Singer 2008) z_1, \dots, z_n are δ -dependent over $\mathbb{K}(x)$ iff $\exists \mathcal{L}_1, \dots, \mathcal{L}_n \in \mathbb{K}[\delta]$, linear δ -operators with coefficients in \mathbb{K} , not all 0, and $g \in \mathbb{K}(x)$:

$$\mathcal{L}_1 \left(\frac{\delta(a_1)}{a_1} \right) + \dots + \mathcal{L}_n \left(\frac{\delta(a_n)}{a_n} \right) = \sigma(g) - g.$$

(Arreche 2017, Arreche-Z. 2022): Using (q) -discrete residues, there exist constants $m_1, \dots, m_n \in \mathbb{K}$, not all 0, such that

$$m_1 \frac{\delta(a_1)}{a_1} + \dots + m_n \frac{\delta(a_n)}{a_n} = \sigma(g) - g + c$$

for some $g \in \mathbb{K}(x)$ and $c \in \mathbb{K}$ (with $c = 0$ in case (S)).

Motivation

Proposition (Hardouin-Singer 2008) z_1, \dots, z_n are δ -dependent over $\mathbb{K}(x)$ iff $\exists \mathcal{L}_1, \dots, \mathcal{L}_n \in \mathbb{K}[\delta]$, linear δ -operators with coefficients in \mathbb{K} , not all 0, and $g \in \mathbb{K}(x)$:

$$\mathcal{L}_1 \left(\frac{\delta(a_1)}{a_1} \right) + \dots + \mathcal{L}_n \left(\frac{\delta(a_n)}{a_n} \right) = \sigma(g) - g.$$

It also holds for the Mahler case. **Question:** How to derive the explicit formulae for $\mathcal{L}_1, \dots, \mathcal{L}_n$?

Motivation

Proposition (Hardouin-Singer 2008) z_1, \dots, z_n are δ -dependent over $\mathbb{K}(x)$ iff $\exists \mathcal{L}_1, \dots, \mathcal{L}_n \in \mathbb{K}[\delta]$, linear δ -operators with coefficients in \mathbb{K} , not all 0, and $g \in \mathbb{K}(x)$:

$$\mathcal{L}_1 \left(\frac{\delta(a_1)}{a_1} \right) + \dots + \mathcal{L}_n \left(\frac{\delta(a_n)}{a_n} \right) = \sigma(g) - g.$$

It also holds for the Mahler case. **Question:** How to derive the explicit formulae for $\mathcal{L}_1, \dots, \mathcal{L}_n$?

Idea: Develop the notion of Mahler discrete residues and derive an effective version of Hardouin-Singer's Proposition in the Mahler case.

Continuous residues

Let \mathbb{K} be an algebraically closed field of char 0, and let $f(x) \in \mathbb{K}(x)$. Make the partial fraction decomposition

$$f(x) = r(x) + \sum_{\alpha \in \mathbb{K}} \sum_{k \geq 1} \frac{c_{\alpha}(k)}{(x - \alpha)^k},$$

where $r(x) \in \mathbb{K}[x]$ and $c_{\alpha}(k) \in \mathbb{K}$ (almost all 0).

Continuous residues

Let \mathbb{K} be an algebraically closed field of char 0, and let $f(x) \in \mathbb{K}(x)$. Make the partial fraction decomposition

$$f(x) = r(x) + \sum_{\alpha \in \mathbb{K}} \sum_{k \geq 1} \frac{c_{\alpha}(k)}{(x - \alpha)^k},$$

where $r(x) \in \mathbb{K}[x]$ and $c_{\alpha}(k) \in \mathbb{K}$ (almost all 0).

Then $f(x)$ is **rationally integrable**, i.e., there exists $g(x) \in \mathbb{K}(x)$ such that $g'(x) = f(x)$, if and only if the (continuous first-order) *residues*

$$\text{res}(f, \alpha, 1) := c_{\alpha}(1) = 0 \quad \text{for every } \alpha \in \mathbb{K}.$$

Continuous residues

Let \mathbb{K} be an algebraically closed field of char 0, and let $f(x) \in \mathbb{K}(x)$. Make the partial fraction decomposition

$$f(x) = r(x) + \sum_{\alpha \in \mathbb{K}} \sum_{k \geq 1} \frac{c_{\alpha}(k)}{(x - \alpha)^k},$$

where $r(x) \in \mathbb{K}[x]$ and $c_{\alpha}(k) \in \mathbb{K}$ (almost all 0).

Then $f(x)$ is **rationally integrable**, i.e., there exists $g(x) \in \mathbb{K}(x)$ such that $g'(x) = f(x)$, if and only if the (continuous first-order) *residues*

$$\text{res}(f, \alpha, 1) := c_{\alpha}(1) = 0 \quad \text{for every } \alpha \in \mathbb{K}.$$

Chen and Singer (2012) created a notion of *discrete residues* that plays an analogous role (where *integrability* \mapsto *summability*) for the *shift* ($x \mapsto x + 1$) and *q-dilation* ($x \mapsto qx$) difference operators.

Discrete residues: shift case

Rewrite the partial fraction decomposition of $f(x) \in \mathbb{K}(x)$:

$$f(x) = r(x) + \sum_{k \geq 1} \sum_{[\alpha] \in \mathbb{K}/\mathbb{Z}} \sum_{n \in \mathbb{Z}} \frac{c_\alpha(k, n)}{(x - \alpha + n)^k}$$

where $r(x) \in \mathbb{K}[x]$, $\alpha \in \mathbb{K}$ is a coset representative for $[\alpha] := \alpha + \mathbb{Z} \in \mathbb{K}/\mathbb{Z}$, and $c_\alpha(k, n) \in \mathbb{K}$.

Discrete residues: shift case

Rewrite the partial fraction decomposition of $f(x) \in \mathbb{K}(x)$:

$$f(x) = r(x) + \sum_{k \geq 1} \sum_{[\alpha] \in \mathbb{K}/\mathbb{Z}} \sum_{n \in \mathbb{Z}} \frac{c_{\alpha}(k, n)}{(x - \alpha + n)^k}$$

where $r(x) \in \mathbb{K}[x]$, $\alpha \in \mathbb{K}$ is a coset representative for $[\alpha] := \alpha + \mathbb{Z} \in \mathbb{K}/\mathbb{Z}$, and $c_{\alpha}(k, n) \in \mathbb{K}$.

The **discrete residue** of $f(x) \in \mathbb{K}(x)$ at the \mathbb{Z} -orbit $[\alpha] \in \mathbb{K}/\mathbb{Z}$ of order k is defined as

$$\text{dres}(f, [\alpha], k) := \sum_{n \in \mathbb{Z}} c_{\alpha}(k, n).$$

Discrete residues: shift case

Rewrite the partial fraction decomposition of $f(x) \in \mathbb{K}(x)$:

$$f(x) = r(x) + \sum_{k \geq 1} \sum_{[\alpha] \in \mathbb{K}/\mathbb{Z}} \sum_{n \in \mathbb{Z}} \frac{c_{\alpha}(k, n)}{(x - \alpha + n)^k}$$

where $r(x) \in \mathbb{K}[x]$, $\alpha \in \mathbb{K}$ is a coset representative for $[\alpha] := \alpha + \mathbb{Z} \in \mathbb{K}/\mathbb{Z}$, and $c_{\alpha}(k, n) \in \mathbb{K}$.

The **discrete residue** of $f(x) \in \mathbb{K}(x)$ at the \mathbb{Z} -orbit $[\alpha] \in \mathbb{K}/\mathbb{Z}$ of order k is defined as

$$\text{dres}(f, [\alpha], k) := \sum_{n \in \mathbb{Z}} c_{\alpha}(k, n).$$

Proposition (Chen-Singer 2012) $f(x)$ is **rationally summable**, i.e., there exists $g(x) \in \mathbb{K}(x)$ such that $f(x) = g(x+1) - g(x)$, if and only if $\text{dres}(f, [\alpha], k) = 0$ for each $[\alpha] \in \mathbb{K}/\mathbb{Z}$ and $k \in \mathbb{N}$.

Discrete residues: q -dilation case

Fix $q \in \mathbb{K}^\times$ not a root of unity, and write $f(x) \in \mathbb{K}(x)$:

$$f(x) = r(x) + \sum_{k \geq 1} \sum_{[\alpha]_q \in \mathbb{K}^\times / q^{\mathbb{Z}}} \sum_{n \in \mathbb{Z}} \frac{c_\alpha(k, n)}{(x - q^n \alpha)^k}$$

where $r(x) = \sum_{j \in \mathbb{Z}} r_j x^j \in \mathbb{K}[x, x^{-1}]$ and $\alpha \in \mathbb{K}^\times$ is a coset representative for $[\alpha]_q := \alpha \cdot q^{\mathbb{Z}} \in \mathbb{K}^\times / q^{\mathbb{Z}}$.

Discrete residues: q -dilation case

Fix $q \in \mathbb{K}^\times$ not a root of unity, and write $f(x) \in \mathbb{K}(x)$:

$$f(x) = r(x) + \sum_{k \geq 1} \sum_{[\alpha]_q \in \mathbb{K}^\times / q^{\mathbb{Z}}} \sum_{n \in \mathbb{Z}} \frac{c_\alpha(k, n)}{(x - q^n \alpha)^k}$$

where $r(x) = \sum_{j \in \mathbb{Z}} r_j x^j \in \mathbb{K}[x, x^{-1}]$ and $\alpha \in \mathbb{K}^\times$ is a coset representative for $[\alpha]_q := \alpha \cdot q^{\mathbb{Z}} \in \mathbb{K}^\times / q^{\mathbb{Z}}$.

The **q -discrete residue** of $f(x)$ at the $q^{\mathbb{Z}}$ -orbit $[\alpha]_q \in \mathbb{K}^\times / q^{\mathbb{Z}}$ of order k (resp., at infinity) is defined as

$$q\text{-dres}(f, [\alpha]_q, k) := \sum_{n \in \mathbb{Z}} q^{-kn} c_\alpha(k, n) \quad (\text{resp., } q\text{-dres}(f, \infty) := r_0).$$

Discrete residues: q -dilation case

Fix $q \in \mathbb{K}^\times$ not a root of unity, and write $f(x) \in \mathbb{K}(x)$:

$$f(x) = r(x) + \sum_{k \geq 1} \sum_{[\alpha]_q \in \mathbb{K}^\times / q^{\mathbb{Z}}} \sum_{n \in \mathbb{Z}} \frac{c_\alpha(k, n)}{(x - q^n \alpha)^k}$$

where $r(x) = \sum_{j \in \mathbb{Z}} r_j x^j \in \mathbb{K}[x, x^{-1}]$ and $\alpha \in \mathbb{K}^\times$ is a coset representative for $[\alpha]_q := \alpha \cdot q^{\mathbb{Z}} \in \mathbb{K}^\times / q^{\mathbb{Z}}$.

The **q -discrete residue** of $f(x)$ at the $q^{\mathbb{Z}}$ -orbit $[\alpha]_q \in \mathbb{K}^\times / q^{\mathbb{Z}}$ of order k (resp., at infinity) is defined as

$$q\text{-dres}(f, [\alpha]_q, k) := \sum_{n \in \mathbb{Z}} q^{-kn} c_\alpha(k, n) \quad (\text{resp., } q\text{-dres}(f, \infty) := r_0).$$

Proposition (Chen-Singer 2012) $f(x)$ is **rationally q -summable**, i.e., there exists $g(x) \in \mathbb{K}(x)$ such that $f(x) = g(qx) - g(x)$, if and only if $q\text{-dres}(f, \infty) = 0$ and $q\text{-dres}(f, [\alpha]_q, k) = 0$ for each $[\alpha]_q \in \mathbb{K}^\times / q^{\mathbb{Z}}$ and $k \in \mathbb{N}$.

Why use residues?

An advantage of using residues is to answer *whether* (yes/no) $f(x) \in \mathbb{K}(x)$ is

- ▶ rationally integrable: $f(x) = g'(x)$; or
- ▶ rationally summable: $f(x) = g(x+1) - g(x)$; or
- ▶ rationally q -summable: $f(x) = g(qx) - g(x)$;

without computing the **certificate** $g(x) \in \mathbb{K}(x)$.

Why use residues?

An advantage of using residues is to answer *whether* (yes/no) $f(x) \in \mathbb{K}(x)$ is

- ▶ rationally integrable: $f(x) = g'(x)$; or
- ▶ rationally summable: $f(x) = g(x+1) - g(x)$; or
- ▶ rationally q -summable: $f(x) = g(qx) - g(x)$;

without computing the **certificate** $g(x) \in \mathbb{K}(x)$.

Computing residues from the definition is impractical because it requires doing partial fraction decompositions.

Why use residues?

An advantage of using residues is to answer *whether* (yes/no) $f(x) \in \mathbb{K}(x)$ is

- ▶ rationally integrable: $f(x) = g'(x)$; or
- ▶ rationally summable: $f(x) = g(x+1) - g(x)$; or
- ▶ rationally q -summable: $f(x) = g(qx) - g(x)$;

without computing the **certificate** $g(x) \in \mathbb{K}(x)$.

Computing residues from the definition is impractical because it requires doing partial fraction decompositions.

In the differential case, there is a better way: if $f = \frac{a}{b}$ with $a, b \in \mathbb{K}[x]$, $\gcd(a, b) = 1$, $\deg(a) < \deg(b)$, and b squarefree, then the roots of the **Rothstein-Trager resultant**

$$RT(f) := \operatorname{Res}_x(a - z \cdot b', b) \in \mathbb{K}[z]$$

are precisely the first-order continuous residues of $f(x)$, which implies $f(x)$ is rationally integrable iff $RT(f)$ is a monomial in z .

Mahler summability for rational functions

Fix $p \in \mathbb{Z}_{\geq 2}$ and let the Mahler difference operator $\sigma : g(x) \mapsto g(x^p)$ for $g(x) \in \mathbb{K}(x)$.

Mahler summability for rational functions

Fix $p \in \mathbb{Z}_{\geq 2}$ and let the **Mahler difference operator**
 $\sigma : g(x) \mapsto g(x^p)$ for $g(x) \in \mathbb{K}(x)$.

We say $f(x) \in \mathbb{K}(x)$ is **Mahler summable** if there exists
 $g(x) \in \mathbb{K}(x)$ such that $f(x) = g(x^p) - g(x)$.

Mahler summability for rational functions

Fix $p \in \mathbb{Z}_{\geq 2}$ and let the **Mahler difference operator** $\sigma : g(x) \mapsto g(x^p)$ for $g(x) \in \mathbb{K}(x)$.

We say $f(x) \in \mathbb{K}(x)$ is **Mahler summable** if there exists $g(x) \in \mathbb{K}(x)$ such that $f(x) = g(x^p) - g(x)$.

Mahler Summability Problem: given $f(x) \in \mathbb{K}(x)$, decide effectively whether $f(x)$ is Mahler summable.

- ▶ Done by Chyzak-Dreyfus-Dumas-Mezzarobba (2018).

Mahler summability for rational functions

Fix $p \in \mathbb{Z}_{\geq 2}$ and let the **Mahler difference operator** $\sigma : g(x) \mapsto g(x^p)$ for $g(x) \in \mathbb{K}(x)$.

We say $f(x) \in \mathbb{K}(x)$ is **Mahler summable** if there exists $g(x) \in \mathbb{K}(x)$ such that $f(x) = g(x^p) - g(x)$.

Mahler Summability Problem: given $f(x) \in \mathbb{K}(x)$, decide effectively whether $f(x)$ is Mahler summable.

► Done by Chyzak-Dreyfus-Dumas-Mezzarobba (2018).

Our Goal: Construct a (\mathbb{K} -linear) complete obstruction to the Mahler summability of $f(x) \in \mathbb{K}(x)$.

Mahler summability for rational functions

More precisely, for the \mathbb{K} -linear map $\Delta : g(x) \mapsto g(x^p) - g(x)$, we wish to construct explicitly a \mathbb{K} -linear map ∇ on $\mathbb{K}(x)$ such that $\text{im}(\Delta) = \ker(\nabla)$, bypassing computation of certificates.

We call ∇ the **Mahler reduction** operator. Given $f \in \mathbb{K}(x)$, set $\bar{f} = \nabla(f)$. Then f is Mahler summable if and only if $\bar{f} = 0$. The numerators in the partial fraction decomposition of \bar{f} are **Mahler discrete residues** of f .

Mahler trajectories and Mahler trees

Let $\mathcal{P} = \{p^n \mid n \in \mathbb{Z}_{\geq 0}\}$ denote the multiplicative monoid of non-negative powers of p .

Mahler trajectories and Mahler trees

Let $\mathcal{P} = \{p^n \mid n \in \mathbb{Z}_{\geq 0}\}$ denote the multiplicative monoid of non-negative powers of p .

We denote by \mathbb{Z}/\mathcal{P} the set of **maximal trajectories** for the action of \mathcal{P} on \mathbb{Z} by multiplication:

$$\mathbb{Z}/\mathcal{P} = \{\{0\}\} \cup \{\{ip^n \mid n \in \mathbb{Z}_{\geq 0}\} \mid i \in \mathbb{Z} \text{ such that } p \nmid i\}.$$

The elements $\theta \in \mathbb{Z}/\mathcal{P}$ are pairwise disjoint subsets of \mathbb{Z} whose union is all of \mathbb{Z} .

Mahler trajectories and Mahler trees

Let $\mathcal{P} = \{p^n \mid n \in \mathbb{Z}_{\geq 0}\}$ denote the multiplicative monoid of non-negative powers of p .

We denote by \mathbb{Z}/\mathcal{P} the set of **maximal trajectories** for the action of \mathcal{P} on \mathbb{Z} by multiplication:

$$\mathbb{Z}/\mathcal{P} = \{\{0\}\} \cup \{\{ip^n \mid n \in \mathbb{Z}_{\geq 0}\} \mid i \in \mathbb{Z} \text{ such that } p \nmid i\}.$$

The elements $\theta \in \mathbb{Z}/\mathcal{P}$ are pairwise disjoint subsets of \mathbb{Z} whose union is all of \mathbb{Z} .

We denote by \mathcal{T}_M the set of equivalence classes for the equivalence relation on \mathbb{K}^\times defined by $\alpha \sim \gamma$ if and only if $\alpha^{p^s} = \gamma^{p^r}$ for some $r, s \in \mathbb{Z}_{\geq 0}$.

The elements $\tau \in \mathcal{T}_M$, called **Mahler trees**, are pairwise disjoint subsets of \mathbb{K}^\times whose union is all of \mathbb{K}^\times . We write $\tau(\alpha)$ for the unique Mahler tree containing $\alpha \in \mathbb{K}^\times$.


Examples of Mahler trees

We define a digraph on the vertex set τ for each Mahler tree $\tau \in \mathcal{T}_M$ with one directed edge $\alpha \rightarrow \gamma$ whenever $\alpha^p = \gamma$.

Examples of Mahler trees

We define a digraph on the vertex set τ for each Mahler tree $\tau \in \mathcal{T}_M$ with one directed edge $\alpha \rightarrow \gamma$ whenever $\alpha^p = \gamma$.


With $p = 3$, the vertices of $\tau(2)$ near $2 \in \mathbb{K}^\times$ are




Examples of Mahler trees

We define a digraph on the vertex set τ for each Mahler tree $\tau \in \mathcal{T}_M$ with one directed edge $\alpha \rightarrow \gamma$ whenever $\alpha^p = \gamma$.

With $p = 3$, the vertices of $\tau(2)$ near $2 \in \mathbb{K}^\times$ are



With $p = 3$, the vertices of $\tau(\zeta_4)$ near $\zeta_4 \in \mathbb{K}^\times$ are



Mahler decomposition of partial fractions

For $f(x) \in \mathbb{K}(x)$, we can decompose it uniquely as

$$f(x) = f_L(x) + f_T(x):$$

$$f_L(x) := \sum_{j \in \mathbb{Z}} r_j x^j \quad \text{and} \quad f_T(x) := \sum_{k \geq 1} \sum_{\alpha \in \mathbb{K}^\times} \frac{c_\alpha(k)}{(x - \alpha)^k},$$

where $r_j, c_\alpha(k) \in \mathbb{K}$.

Mahler decomposition of partial fractions

For $f(x) \in \mathbb{K}(x)$, we can decompose it uniquely as

$$f(x) = f_L(x) + f_T(x):$$

$$f_L(x) := \sum_{j \in \mathbb{Z}} r_j x^j \quad \text{and} \quad f_T(x) := \sum_{k \geq 1} \sum_{\alpha \in \mathbb{K}^\times} \frac{c_\alpha(k)}{(x - \alpha)^k},$$

where $r_j, c_\alpha(k) \in \mathbb{K}$.

This yields a σ -stable \mathbb{K} -vector space decomposition of $\mathbb{K}(x)$ as

$$\mathbb{K}(x) \simeq \mathbb{K}[x, x^{-1}] \oplus \mathbb{K}(x)_T.$$

Mahler decomposition of partial fractions

For $f(x) \in \mathbb{K}(x)$, we can decompose it uniquely as

$$f(x) = f_L(x) + f_T(x):$$

$$f_L(x) := \sum_{j \in \mathbb{Z}} r_j x^j \quad \text{and} \quad f_T(x) := \sum_{k \geq 1} \sum_{\alpha \in \mathbb{K}^\times} \frac{c_\alpha(k)}{(x - \alpha)^k},$$

where $r_j, c_\alpha(k) \in \mathbb{K}$.

This yields a σ -stable \mathbb{K} -vector space decomposition of $\mathbb{K}(x)$ as

$$\mathbb{K}(x) \simeq \mathbb{K}[x, x^{-1}] \oplus \mathbb{K}(x)_T.$$

Then $f \in \mathbb{K}(x)$ is Mahler summable if and only if f_L and f_T are both Mahler summable. We address each component separately.

Mahler decomposition of partial fractions

For $f(x) \in \mathbb{K}(x)$, we can decompose it uniquely as

$f(x) = f_L(x) + f_T(x)$:

$$f_L(x) := \sum_{j \in \mathbb{Z}} r_j x^j \quad \text{and} \quad f_T(x) := \sum_{k \geq 1} \sum_{\alpha \in \mathbb{K}^\times} \frac{c_\alpha(k)}{(x - \alpha)^k},$$

where $r_j, c_\alpha(k) \in \mathbb{K}$.

This yields a σ -stable \mathbb{K} -vector space decomposition of $\mathbb{K}(x)$ as

$$\mathbb{K}(x) \simeq \mathbb{K}[x, x^{-1}] \oplus \mathbb{K}(x)_T.$$

Then $f \in \mathbb{K}(x)$ is Mahler summable if and only if f_L and f_T are both Mahler summable. We address each component separately.

Moreover, the decompositions $f_L = \sum_{\theta \in \mathbb{Z}/\mathcal{P}} f_\theta$ and $f_T = \sum_{\tau \in \mathcal{T}_M} f_\tau$:

$$f_\theta := \sum_{j \in \theta} r_j x^j \quad \text{and} \quad f_\tau := \sum_{k \geq 1} \sum_{\alpha \in \tau} \frac{c_\alpha(k)}{(x - \alpha)^k}$$

are also σ -stable. Can decide summability of f by deciding for each f_θ ($\theta \in \mathbb{Z}/\mathcal{P}$) and each f_τ ($\tau \in \mathcal{T}_M$) individually.

Mahler residues at infinity

Definition (Arreche-Z. 2022) Let $f(x) \in \mathbb{K}(x)$ and write $f_L(x) = \sum_{j \in \mathbb{Z}} r_j x^j \in \mathbb{K}[x, x^{-1}]$. The **Mahler residue** of $f(x)$ at infinity is the vector

$$\text{dres}(f, \infty) := \left(\sum_{j \in \theta} r_j \right)_{\theta \in \mathbb{Z}/\mathcal{P}} \in \bigoplus_{\theta \in \mathbb{Z}/\mathcal{P}} \mathbb{K}.$$

Mahler residues at infinity

Definition (Arreche-Z. 2022) Let $f(x) \in \mathbb{K}(x)$ and write $f_L(x) = \sum_{j \in \mathbb{Z}} r_j x^j \in \mathbb{K}[x, x^{-1}]$. The **Mahler residue** of $f(x)$ at infinity is the vector

$$\text{dres}(f, \infty) := \left(\sum_{j \in \theta} r_j \right)_{\theta \in \mathbb{Z}/\mathcal{P}} \in \bigoplus_{\theta \in \mathbb{Z}/\mathcal{P}} \mathbb{K}.$$

Proposition (Arreche-Z. 2022) For $f(x) \in \mathbb{K}(x)$ the component $f_L(x) \in \mathbb{K}[x, x^{-1}]$ is Mahler summable if and only if $\text{dres}(f, \infty) = \mathbf{0}$ (the zero vector).

Mahler residues at Mahler trees

Suppose $\gamma \in \mathbb{K}^\times$ is not a root of unity, $f \in \mathbb{K}(x)$, and $h \in \mathbb{Z}_{\geq 0}$ s.t.:

$$\text{sing}(f) \cap \tau(\gamma) \subseteq \{\zeta_{p^n}^i \gamma^{p^{h-n}} \mid 0 \leq n \leq h, i \in \mathbb{Z}/p^n\mathbb{Z}\}.$$

Mahler residues at Mahler trees

Suppose $\gamma \in \mathbb{K}^\times$ is not a root of unity, $f \in \mathbb{K}(x)$, and $h \in \mathbb{Z}_{\geq 0}$ s.t.:

$$\text{sing}(f) \cap \tau(\gamma) \subseteq \{\zeta_{p^n}^i \gamma^{p^{h-n}} \mid 0 \leq n \leq h, i \in \mathbb{Z}/p^n\mathbb{Z}\}.$$

Then we can write, for $\tau = \tau(\gamma)$,

$$f_\tau = \sum_{k=1}^m \sum_{n=0}^h \sum_{i \in \mathbb{Z}/p^n\mathbb{Z}} \frac{c_\gamma(k, n, i)}{(x - \zeta_{p^n}^i \gamma^{p^{h-n}})^k}.$$

Mahler residues at Mahler trees

Suppose $\gamma \in \mathbb{K}^\times$ is not a root of unity, $f \in \mathbb{K}(x)$, and $h \in \mathbb{Z}_{\geq 0}$ s.t.:

$$\text{sing}(f) \cap \tau(\gamma) \subseteq \{\zeta_{p^n}^i \gamma^{p^{h-n}} \mid 0 \leq n \leq h, i \in \mathbb{Z}/p^n\mathbb{Z}\}.$$

Then we can write, for $\tau = \tau(\gamma)$,

$$f_\tau = \sum_{k=1}^m \sum_{n=0}^h \sum_{i \in \mathbb{Z}/p^n\mathbb{Z}} \frac{c_\gamma(k, n, i)}{(x - \zeta_{p^n}^i \gamma^{p^{h-n}})^k}.$$

Set recursively: $\tilde{c}_{k,0,0} := c_\gamma(k, 0, 0)$, and for $1 \leq n \leq h$; $i \in \mathbb{Z}/p^n\mathbb{Z}$:

$$\tilde{c}_{k,n,i} := c_\gamma(k, n, i) + \sum_{j=k}^m \tilde{c}_{j,n-1,\pi_{n-1}(i)} V_k^j(\zeta_{p^n}^i \gamma^{p^{h-n}}),$$

where $\pi_{n-1} : \mathbb{Z}/p^n\mathbb{Z} \rightarrow \mathbb{Z}/p^{n-1}\mathbb{Z}$ is the canonical projection.

Mahler residues at Mahler trees

Suppose $\gamma \in \mathbb{K}^\times$ is not a root of unity, $f \in \mathbb{K}(x)$, and $h \in \mathbb{Z}_{\geq 0}$ s.t.:

$$\text{sing}(f) \cap \tau(\gamma) \subseteq \{\zeta_{p^n}^i \gamma^{p^{h-n}} \mid 0 \leq n \leq h, i \in \mathbb{Z}/p^n\mathbb{Z}\}.$$

Then we can write, for $\tau = \tau(\gamma)$,

$$f_\tau = \sum_{k=1}^m \sum_{n=0}^h \sum_{i \in \mathbb{Z}/p^n\mathbb{Z}} \frac{c_\gamma(k, n, i)}{(x - \zeta_{p^n}^i \gamma^{p^{h-n}})^k}.$$

Set **recursively**: $\tilde{c}_{k,0,0} := c_\gamma(k, 0, 0)$, and for $1 \leq n \leq h$; $i \in \mathbb{Z}/p^n\mathbb{Z}$:

$$\tilde{c}_{k,n,i} := c_\gamma(k, n, i) + \sum_{j=k}^m \tilde{c}_{j,n-1,\pi_{n-1}(i)} V_k^j(\zeta_{p^n}^i \gamma^{p^{h-n}}),$$

where $\pi_{n-1} : \mathbb{Z}/p^n\mathbb{Z} \rightarrow \mathbb{Z}/p^{n-1}\mathbb{Z}$ is the canonical projection.

Mahler residues at Mahler trees

Suppose $\gamma \in \mathbb{K}^\times$ is not a root of unity, $f \in \mathbb{K}(x)$, and $h \in \mathbb{Z}_{\geq 0}$ s.t.:

$$\text{sing}(f) \cap \tau(\gamma) \subseteq \{\zeta_{p^n}^i \gamma^{p^{h-n}} \mid 0 \leq n \leq h, i \in \mathbb{Z}/p^n\mathbb{Z}\}.$$

Then we can write, for $\tau = \tau(\gamma)$,

$$f_\tau = \sum_{k=1}^m \sum_{n=0}^h \sum_{i \in \mathbb{Z}/p^n\mathbb{Z}} \frac{c_\gamma(k, n, i)}{(x - \zeta_{p^n}^i \gamma^{p^{h-n}})^k}.$$

Set recursively: $\tilde{c}_{k,0,0} := c_\gamma(k, 0, 0)$, and for $1 \leq n \leq h$; $i \in \mathbb{Z}/p^n\mathbb{Z}$:

$$\tilde{c}_{k,n,i} := c_\gamma(k, n, i) + \sum_{j=k}^m \tilde{c}_{j,n-1,\pi_{n-1}(i)} V_k^j(\zeta_{p^n}^i \gamma^{p^{h-n}}),$$

where $\pi_{n-1} : \mathbb{Z}/p^n\mathbb{Z} \rightarrow \mathbb{Z}/p^{n-1}\mathbb{Z}$ is the canonical projection.

Definition (Arreche-Z. 2022) The **Mahler discrete residue** at τ of order k is the vector $\text{dres}(f, \tau, k) \in \bigoplus_{\alpha \in \tau} \mathbb{K}$ with α -component $:= 0$ except possibly at $\alpha = \zeta_{p^h}^i \gamma$ for $i \in \mathbb{Z}/p^h\mathbb{Z}$, given by $\tilde{c}_{k,h,i}$.

Mahler residues at Mahler trees

Proposition (Arreche-Z. 2022) For $f \in \mathbb{K}(x)$ the component $f_{\mathcal{T}}$ is Mahler summable if and only if $\text{dres}(f, \tau, k) = \mathbf{0}$ (the zero vector) for each $\tau \in \mathcal{T}_M$ and $k \in \mathbb{N}$.

Mahler residues at Mahler trees

Proposition (Arreche-Z. 2022) For $f \in \mathbb{K}(x)$ the component $f_{\mathcal{T}}$ is Mahler summable if and only if $\text{dres}(f, \tau, k) = \mathbf{0}$ (the zero vector) for each $\tau \in \mathcal{T}_M$ and $k \in \mathbb{N}$.

Mahler residues at Mahler trees

Proposition (Arreche-Z. 2022) For $f \in \mathbb{K}(x)$ the component f_{τ} is Mahler summable if and only if $\text{dres}(f, \tau, k) = \mathbf{0}$ (the zero vector) for each $\tau \in \mathcal{T}_M$ and $k \in \mathbb{N}$.

- ▶ The definition (and proofs) for Mahler discrete residues at $\tau(\zeta)$ for $\zeta \in \mathbb{K}_t^{\times}$ a root of unity is similar in spirit, but more technical, due to the perverse (pre-)periodic behavior of roots of unity under the p -power map.

Main Result

Theorem (Arreche-Z. 2022) Given $f \in \mathbb{K}(x)$. Then f is Mahler summable if and only if $\text{dres}(f, \infty) = \mathbf{0}$ and $\text{dres}(f, \tau, k) = \mathbf{0}$ for all $k \in \mathbb{N}$ and $\tau \in \mathcal{T}_M$.

Thanks!