

# On Sequences Associated to the Invariant Theory of Rank Two Lie Algebras

Yi Zhang

Department of Foundational Mathematics  
Xi'an Jiaotong-Liverpool University, Suzhou, China

Joint work with Alin Bostan, Jordan Tirrell and Bruce Westbury





## Two families of sequences in OEIS

a (OEIS tag)	0	1	2	3	4	5
A059710	1	0	1	1	4	10
A108307	1	1	2	5	15	51
A108304	1	2	5	15	52	202

The first family of sequences (**octant sequences**)

a (OEIS tag)	0	1	2	3	4	5
A151366	1	0	2	2	12	30
A236408	1	1	3	9	33	131
A001181	1	2	6	22	92	422
A216947	1	3	11	49	221	1113

The second family of sequences (**quadrant sequences**)

- ▶ Those sequences are associated to the invariant theory of the exceptional simple Lie algebra  $G_2$  of rank 2.
- ▶ The quadrant sequences are related to the octant sequences by the branching rules for  $SL(3)$  of  $G_2$ .

# Octant sequences

The first family of sequences can be interpreted as lattice walks restricted to the octant. We call them **octant sequences**.

- ▶ **A059710**: enumerates the multiplicities of the trivial representation in the tensor powers of  $V$ , which is the 7-D fundamental representation of  $G_2$ .
- ▶ **A108307**: enumerates **enhanced** 3-noncrossing set partitions.
- ▶ **A108304**: enumerates 3-noncrossing set partitions.

(Lin, 2018; Gil and Tirrell, 2019): **A108307** and **A108304** are related by the binomial transform.

## Motivation and Contribution

(Bostan, Tirrell, Westbury and Z., 2019): [A059710](#) and [A108307](#) are also related by the binomial transform.

**Mihailovs' conjecture:** Let  $T_3(n)$  be the  $n$ -th term of [A059710](#). Then  $T_3$  is determined by  $T_3(0) = 1$ ,  $T_3(1) = 0$ ,  $T_3(2) = 1$  and

$$14(n+1)(n+2)T_3(n) + (n+2)(19n+75)T_3(n+1) + 2(n+2)(2n+11)T_3(n+2) - (n+8)(n+9)T_3(n+3) = 0.$$

(Bostan, Tirrell, Westbury and Z., 2019): Three independent proofs of Mihailovs' conjecture.

- ▶ Two proofs are based on binomial relation between [A059710](#) and [A108307](#), together with a result by Bousquet-Mélou and Xin.
- ▶ The third one is a direct proof by the method of algebraic residues, which leads to closed formulae for the generating function of  $T_3$  in terms of hypergeometric functions.

## Quadrant sequences

The second family of sequences can be interpreted as lattice walks restricted to the quadrant. We call them **quadrant sequences**.

- ▶ [A151366](#): enumerates nonpositive bipartite trivalent graphs.
- ▶ [A236408](#): enumerates pasting diagrams.
- ▶ [A001181](#): enumerates Baxter permutations.
- ▶ [A216947](#): enumerates 2-coloured noncrossing set partitions.

**Question:** What are relations between quadrant sequences?

# Motivation and Contribution

(Marberg, 2013): a combinatorial proof that [A151366](#), [A001181](#), and [A216947](#) are related by binomial transforms.

(Bostan, Tirrell, Westbury and Z., 2019): Derive a uniform recurrence equation for quadrant sequences and show that they are related by binomial transform by the representation theory of simple Lie algebras.

# Outline

- ▶ binomial relation between the first and second octant sequences
- ▶ Three independent proofs of Mihailovs' conjecture
- ▶ Recurrence relations for the quadrant sequences



# Preliminaries

**Definition 1** Let  $G$  be a reductive complex algebraic group and let  $V$  be a representation of  $G$ . The sequence associated to  $(G, V)$ , denoted  $\mathbf{a}_V$ , is the sequence whose  $n$ -th term is the multiplicity of the trivial representation in the tensor power  $\otimes^n V$ .

**Example 1** Let  $V$  be the 7-D fundamental representation of  $G_2$ . Then [A059710](#) is the sequence associated with  $(G_2, V)$ .

Let  $\mathbf{a}$  be a sequence with  $n$ -th term  $a(n)$ , the **binomial transform** of  $\mathbf{a}$  is the sequence, denoted  $\mathcal{B}\mathbf{a}$ , whose  $n$ -th term is

$$\sum_{i=0}^n \binom{n}{i} a(i).$$

## Preliminaries

**Lemma 1** Assume  $\mathbf{a}_V$  is the sequence associated to  $(G, V)$  as specified in **Definition 1**. Then  $\mathbf{a}_{V \oplus \mathbb{C}} = \mathcal{B}\mathbf{a}_V$ .

**Lemma 2** Assume  $\mathbf{a}$  enumerates walks in a lattice, confined to a domain  $D$ , using a set of steps  $S$ . Then  $\mathcal{B}\mathbf{a}$  also enumerates walks in a lattice restricted to  $D$  with steps  $S \amalg \{0\}$ .

**Lemma 3** Let  $G(t)$  be the generating function of  $\mathbf{a}$ . For  $k \in \mathbb{Z}$ , denote the generating function of  $\mathcal{B}^k \mathbf{a}$  by  $\mathcal{B}^k G$ . Then


$$(\mathcal{B}^k G)(t) = \frac{1}{1 - k t} G\left(\frac{t}{1 - k t}\right).$$

## Binomial relation between A059710 and A108307


Let  $V$  be the 7-D fundamental representation of  $G_2$ . Then

- ▶ A059710 is the sequence associated to  $(G_2, V)$ . Let  $T_3(n)$  be its  $n$ -th term.
- ▶ A108307 enumerates enhanced 3-noncrossing set partitions. Let  $E_3(n)$  be its  $n$ -th term.

In terms of lattice walks, we can interpret  $T_3$  and  $E_3$  as follows:




Steps in weight  
lattice of  $G_2$




Steps in octant  
related to  $E_3(n)$

In terms of lattice walks, we can interpret  $T_3$  and  $E_3$  as follows:



Steps in weight  
lattice of  $G_2$



Steps in octant  
related to  $E_3(n)$

If we make a linear transformation  $(x, y) \rightarrow (x + y, y)$ , then it identifies the six non-zero steps, as well as the two domains.

## Binomial relation between A059710 and A108307

**Recall:** **Lemma 2** Assume  $\mathbf{a}$  enumerates walks in a lattice, confined to a domain  $D$ , using a set of steps  $S$ . Then  $\mathcal{B}\mathbf{a}$  also enumerates walks in a lattice restricted to  $D$  with steps  $S \amalg \{0\}$ .

## Binomial relation between A059710 and A108307

**Recall:** [Lemma 2](#) Assume  $\mathbf{a}$  enumerates walks in a lattice, confined to a domain  $D$ , using a set of steps  $S$ . Then  $\mathcal{B}\mathbf{a}$  also enumerates walks in a lattice restricted to  $D$  with steps  $S \amalg \{0\}$ .

By [Lemma 2](#) and the previous figures, we conclude that  $E_3$  is the binomial transform of  $T_3$ .

## Binomial relation between A059710 and A108307

**Recall:** Lemma 2 Assume  $\mathbf{a}$  enumerates walks in a lattice, confined to a domain  $D$ , using a set of steps  $S$ . Then  $\mathcal{B}\mathbf{a}$  also enumerates walks in a lattice restricted to  $D$  with steps  $S \amalg \{0\}$ .

By Lemma 2 and the previous figures, we conclude that  $E_3$  is the binomial transform of  $T_3$ .

(Lin, 2018; Gil and Tirrell, 2019): A108307 and A108304 are related by the binomial transform.

**Recall:** Lemma 1 Assume  $\mathbf{a}_V$  is the sequence associated to  $(G, V)$  as specified in Definition 1. Then  $\mathbf{a}_{V \oplus \mathbb{C}} = \mathcal{B}\mathbf{a}_V$ .

Thus, the octant sequences are sequences associated to

$$(G_2, V), \quad (G_2, V \oplus \mathbb{C}), \quad (G_2, V \oplus 2\mathbb{C}).$$



## First proof of Mihailovs' conjecture

**Mihailovs' conjecture:** Let  $T_3(n)$  be the  $n$ -th term of [A059710](#). Then  $T_3$  is determined by  $T_3(0) = 1$ ,  $T_3(1) = 0$ ,  $T_3(2) = 1$  and

$$14(n+1)(n+2)T_3(n) + (n+2)(19n+75)T_3(n+1) + 2(n+2)(2n+11)T_3(n+2) - (n+8)(n+9)T_3(n+3) = 0.$$

([Bousquet-Mélou and Xin, 2005](#)): Let  $E_3(n)$  be the  $n$ -th term of [A108307](#). Then  $E_3$  is given by  $E_3(0) = E_3(1) = 1$ , and

$$8(n+3)(n+1)E_3(n) + (7n^2 + 53n + 88)E_3(n+1) - (n+8)(n+7)E_3(n+2) = 0.$$

## First proof of Mihailovs' conjecture

**Recall:** We prove that  $E_3$  is the binomial transform of  $T_3$ . Thus,

$$T_3(n) = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} E_3(k).$$

Set  $f(n, k) = (-1)^{n-k} \binom{n}{k} E_3(k)$ .

- ▶ By Bousquet-Mélou and Xin's result,  $f(n, k)$  is holonomic function, which satisfies ordinary difference equations for  $n$  and  $k$ , respectively.
- ▶ **Idea:** Using creative telescoping method (Zeilberger, 1990), which is an algorithmic approach to compute a differential/difference equation for the integration/sum of holonomic functions, to drive a recurrence equation for  $T_3$ .

# First proof of Mihailovs' conjecture

- ▶ Using the Koutschan's Mathematica package `HolonomicFunctions.m` that implements Chyzak's algorithm for creative telescoping, we find exactly the recurrence equation in Mihailovs' conjecture.

## Second proof of Mihailovs' conjecture

**Recall:** We prove that  $E_3$  is the binomial transform of  $T_3$ . Let  $\mathcal{T}(t) = \sum_{n \geq 0} T_3(n)t^n$  and  $\mathcal{E}(t) = \sum_{n \geq 0} E_3(n)t^n$ . Then

$$\mathcal{T}(t) = \frac{1}{1+t} \cdot \mathcal{E}\left(\frac{t}{1+t}\right).$$

- ▶ By Bousquet-Mélou and Xin's result, we can derive an ODE for  $\mathcal{E}(t)$ .
- ▶ Using the closure properties of holonomic function (the sum, product and algebraic substitution of holonomic functions is still holonomic), we can derive an ODE for  $\mathcal{T}(t)$  and convert it into a linear recurrence for  $T_3(n)$ , which is exactly the recurrence equation in Mihailovs' conjecture.

## Third proof of Mihailovs' conjecture

**Idea:** In terms of lattice walks, we can interpret  $T_3(n)$  to be the constant term of  $W K^n$ , where

$$K = (1 + x + y + xy + x^{-1} + y^{-1} + (xy)^{-1}),$$

and

$$W = x^{-2}y^{-3}(x^2y^3 - xy^3 + x^{-1}y^2 - x^{-2}y + x^{-3}y^{-1} - x^{-3}y^{-2} + x^{-2}y^{-3} - x^{-1}y^{-3} + xy^{-2} - x^2y^{-1} + x^3y - x^3y^2).$$

Let  $\mathcal{T}(t) = \sum_{n \geq 0} T_3(n)t^n$ . Then  $\mathcal{T}(t)$  is the constant coefficient  $[x^0y^0]$  of  $W/(1 - tK)$ . In other words,  $\mathcal{T}(t)$  is equal to the algebraic residue of  $W/(xy - txyK)$ , which is proportional to the contour integral of  $W/(xy - txyK)$  over a cycle.

## Third proof of Mihailovs' conjecture

Using creative telescoping method, we can compute a 6-th order ODE for  $\mathcal{T}(t)$ . Moreover, by using factorization of differential operators, we can show that  $L_3(\mathcal{T}(t)) = 0$ , where  $\partial = \frac{d}{dt}$  and

$$L_3 = t^2 (2t + 1) (7t - 1) (t + 1) \partial^3 + 2t(t + 1) (63t^2 + 22t - 7) \partial^2 + (252t^3 + 338t^2 + 36t - 42) \partial + 28t(3t + 4).$$

Converting it into a linear recurrence for  $T_3(n)$ , we get exactly the recurrence equation in Mihailovs' conjecture.

## Closed formulae

By factorization of the operator  $L_3$  and algorithms for solving 2-nd order ODEs, we derive the following closed formula for  $\mathcal{T}(t)$ :

$$\mathcal{T}(t) = \frac{1}{30 t^5} \left[ R_1 \cdot {}_2F_1 \left( \frac{1}{3}, \frac{2}{3}; \phi \right) + R_2 \cdot {}_2F_1 \left( \frac{2}{3}, \frac{4}{3}; \phi \right) + 5 P \right],$$

where

$$R_1 = \frac{(t+1)^2 (214 t^3 + 45 t^2 + 60 t + 5)}{t-1},$$

$$R_2 = 6 \frac{t^2 (t+1)^2 (101 t^2 + 74 t + 5)}{(t-1)^2},$$

and

$$\phi = \frac{27(t+1)t^2}{(1-t)^3}, \quad P = 28 t^4 + 66 t^3 + 46 t^2 + 15 t + 1.$$

## Closed formulae

By elliptic curve theory, we derive an alternative formula for  $\mathcal{T}(t)$ :

$$\frac{P}{6t^5} + \frac{(7t-1)(2t+1)(t+1)}{360t^5} \left( (155t^2 + 182t + 59)(11t+1)H(t) + (341t^3 + 507t^2 + 231t + 1)(5t+1)H'(t) \right),$$

where

$$H(t) = \frac{1}{g_2^{1/4}} \cdot {}_2F_1 \left( \begin{matrix} \frac{1}{12} & \frac{5}{12} \\ 1 \end{matrix}; \frac{1728}{J} \right),$$
$$J = \frac{(t-1)^3 (25t^3 + 21t^2 + 3t - 1)^3}{t^6 (1-7t)(2t+1)^2 (t+1)^3},$$

and

$$g_2 = (t-1)(25t^3 + 21t^2 + 3t - 1).$$



## Transcendence and asymptotics

Using those closed formulae, we can show that that  $\mathcal{T}(t)$  is a transcendental power series and its  $n$ -th coefficient

$$T_3(n) \sim C \cdot \frac{7^n}{n}, \quad \text{where } C = \frac{4117715 \sqrt{3}}{864 \pi} \approx 2627.6.$$

## Recurrence relations for quadrant sequences

**Definition 2** Let  $\tilde{V}$  be the defining representation of  $SL(3)$  and denote the dual by  $\tilde{V}^*$ . For  $k \geq 0$ , we define  $\mathcal{S}_k$  to be the sequence associated to  $(SL(3), \tilde{V} \oplus \tilde{V}^* \oplus k\mathbb{C})$ .

**Remark:**  $SL(3)$  is the maximal subgroup of  $G_2$ . Let  $V$  be the 7-D fundamental representation of  $G_2$ . Then  $\mathcal{S}_k$  is the the sequence associated to  $(SL(3), (V \oplus k\mathbb{C}) \downarrow_{SL(3)})$ .

**Theorem (Bostan, Tirrell, Westbury and Z., 2019):** The quadrant sequences  $\mathcal{S}_0, \mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3$  are identical to the sequences in the second family listed in OEIS.

**Lemma 4** Let  $\mathcal{G}_k$  be the generating function of  $\mathcal{S}_k$ , where  $k \geq 0$ . Then  $\mathcal{G}_k$  is the constant coefficient of  $[x^0 y^0]$  of  $W/(1 - tK)$ , where

$$K = k + x + y + x^{-1} + y^{-1} + \frac{x}{y} + \frac{y}{x}$$

and

$$W = 1 - \frac{x^2}{y} + x^3 - x^2 y^2 + y^3 - \frac{y^2}{x}.$$

## Recurrence relations for quadrant sequences

By [Lemma 4](#),  $\mathcal{S}_3$  is identical to the sequence [A216947](#).

([Marberg, 2013](#)): The  $n$ -th term  $C_2(n)$  of  $\mathcal{S}_3$  is given by  $C_2(0) = 1$ ,  $C_2(1) = 3$  and

$$(n+5)(n+6) \cdot C_2(n+2) - 2(5n^2 + 36n + 61) \cdot C_2(n+1) + 9(n+1)(n+4) \cdot C_2(n) = 0.$$

By [Lemma 1](#),  $\mathcal{S}_k$ 's are related by binomial transforms. Thus, by [Lemma 3](#), the generating function of  $\mathcal{S}_k$  is

$$\mathcal{G}_k(t) = \frac{1}{1-kt} \cdot \mathcal{G}_3\left(\frac{t}{1-kt}\right)$$

where  $\mathcal{G}_3(t)$  is the generating function of  $\mathcal{S}_3$ .

## Recurrence relations for quadrant sequences

Using closure properties of holonomic functions, we derive a uniform 4-th order recurrence equation for  $\mathcal{S}_k$  with  $k$  as a parameter.

By comparing the recurrence equations between  $\mathcal{S}_k$ 's and the sequences in the second family, and then checking initial terms, we show that

**Corollary:** The recurrence relations stated in OEIS for the sequences in the second family are true.

# Summary

- ▶ A combinatorial proof of binomial relation between the first and second octant sequences
- ▶ Three independent proofs of Mihailovs' conjecture
  - ▶ Two proofs are based on binomial relation between the first and second octant sequences
  - ▶ A direct proof by the method of algebraic residues, which leads to closed formulae for the generating function of the first octant sequence
- ▶ A unified proof for recurrence relations of the quadrant sequences

## Summary

- ▶ A combinatorial proof of binomial relation between the first and second octant sequences
- ▶ Three independent proofs of Mihailovs' conjecture
  - ▶ Two proofs are based on binomial relation between the first and second octant sequences
  - ▶ A direct proof by the method of algebraic residues, which leads to closed formulae for the generating function of the first octant sequence
- ▶ A unified proof for recurrence relations of the quadrant sequences

Thanks!