

# Rational Solutions of First-Order Algebraic Ordinary Difference Equations

Yi Zhang

Department of Foundational Mathematics  
Xi'an Jiaotong-Liverpool University, Suzhou, China

Joint work with Thieu N. Vo



# AO $\Delta$ E

Let  $\mathbb{K}$  be an algebraically closed field of char 0, and  $x$  be an indeterminate.

Consider the algebraic ordinary difference equation (AO $\Delta$ E):

$$F(x, y(x), y(x+1), \dots, y(x+m)) = 0, \quad (1)$$

where  $F$  is a polynomial in  $y(x), y(x+1), \dots, y(x+m)$  with coeffs in  $\mathbb{K}(x)$  and  $m \in \mathbb{N}$  is called the **order** of  $F$ . We also simply write (1) as  $F(y) = 0$ . An AO $\Delta$ E is **autonomous** if  $x$  does not appear in it explicitly.

**Example 1.** Equations of Riccati type:

$$y(x+1)y(x) + p(x)y(x+1) + q(x)y(x) = 0,$$

where  $p, q \in \mathbb{K}[x]$ .

# Motivation

**Goal:** Given a first-order AODE  $F(y) = 0$ . Determine a **strong rational general solution**  $s \in \mathbb{K}(x, c) \setminus \mathbb{K}(x)$ , where  $c$  is transcendental over  $\mathbb{K}(x)$ , s.t.

$$F(x, s(x), s(x+1)) = 0.$$

Let  $s(x) = \frac{p(x)}{q(x)}$  with  $\gcd(p, q) = 1$ . Denote the degree of  $s$  by  $\deg(s) := \max(\deg(p), \deg(q))$ .

## Applications:

- ▶ Automatic proof of combinatorial identities: symbolic summation.
- ▶ Difference Galois theory: factorization of linear difference operators.
- ▶ Analysis of time or space complexity of computer programs.

# Motivation

## Previous works:

- ▶ (Abramov-Bronstein-Petkovšek-van Hoeij 1989-1998): Algorithms for computing rational solutions of **linear** difference equations.
- ▶ (Feng-Gao-Huang 2008): An algorithm for computing rational solutions of first-order autonomous AO $\Delta$ Es **provided the degree of the rational solution is given.**
- ▶ (Shkaravska-Eekelen 2014, 2021): a degree bound for polynomial solutions of high-order non-autonomous AO $\Delta$ Es under a sufficient condition.

**Our contribution:** Construct a degree bound for rational solutions of first-order autonomous AO $\Delta$ Es, thus derive a complete algorithm for computing corresponding rational solutions.

# Preliminaries

Let  $F \in \mathbb{K}[x, y, z] \setminus \{0\}$  be an irreducible polynomial.

**Theorem 1:** If the AODE  $F(x, y(x), y(x+1)) = 0$  admits a strong rational general solution, then the algebraic curve in  $\mathbb{A}^2(\overline{\mathbb{K}(x)})$  defined by  $F(x, y, z) = 0$  is of genus zero.

# Preliminaries

Using parametrization theory of rational curves, we have

**Proposition 1:** If the algebraic curve  $\mathcal{C}_F \subset \mathbb{A}^2(\overline{\mathbb{K}(x)})$  defined by  $F(x, y, z) = 0$  is of genus zero, then there exists a birational transformation  $\mathcal{P} : \mathbb{A}^1(\overline{\mathbb{K}(x)}) \rightarrow \mathcal{C}_F$  defined by  $\mathcal{P}(x, t) = (p_1(x, t), p_2(x, t))$  for some  $p_1(x, t), p_2(x, t) \in \mathbb{K}(x, t)$ .

- ▶ There exists an algorithm (Vo-Grasegger-Winkler 2018) for determining such a birational transformation as above.

## Preliminaries

**Theorem 2:** Let  $F(x, y(x), y(x+1)) = 0$  be an AO $\Delta$ E s.t. its corresponding curve  $\mathcal{C}_F$  is of genus zero. Assume  $\mathcal{P}(x, t) = (p_1(x, t), p_2(x, t)) \in \mathbb{K}(x, t)^2$  is a birational transformation from  $\mathbb{A}^1(\overline{\mathbb{K}(x)})$  to  $\mathcal{C}_F$ . Consider

$$p_1(x+1, \omega(x+1)) = p_2(x, \omega(x)). \quad (2)$$

- ▶ If  $s(x, c)$  is a strong rational general solution of  $F(y) = 0$ , then there exists a strong rational general solution  $\omega(x, c)$  of (2) s.t.  $s(x, c) = p_1(x, \omega(x, c))$ .
- ▶ Conversely, if  $\omega(x, c)$  is a strong rational general solution of (2), then  $s(x, c) = p_1(x, \omega(x, c))$  is a strong rational general solution of  $F(y) = 0$ .

We call (2) an **associated separable AO $\Delta$ E** of  $F(y) = 0$ .

## Preliminaries

**Proposition 2:** If the AO $\Delta$ E  $F(x, y(x), y(x+1)) = 0$  admits a strong rational general solution, then we have

$$\deg_y F = \deg_z F.$$

In this case, the associated separable AO $\Delta$ E exists and it must be of the form

$$P(x, \omega(x+1)) = Q(x, \omega(x)),$$

for some  $P, Q \in \mathbb{K}(x, y)$  s.t.

$$\deg_y P = \deg_y Q = \deg_z F = \deg_y F.$$

**Goal:** Construct a degree bound for rational solutions of autonomous separable AO $\Delta$ Es, and thus derive an algorithm for computing rational solutions of first-order autonomous AO $\Delta$ Es.



# Problem

**Question 1:** Let  $P_1, P_2, Q_1, Q_2$  be polynomials in  $\mathbb{K}[z] \setminus \{0\}$  such that  $\gcd(P_1, Q_1) = \gcd(P_2, Q_2) = 1$  and  $\deg \frac{P_1}{Q_1} = \deg \frac{P_2}{Q_2} = n \geq 1$ . Find all rational solutions of the autonomous separable AODE

$$\frac{P_1(y(x+1))}{Q_1(y(x+1))} = \frac{P_2(y(x))}{Q_2(y(x))}. \quad (3)$$

# Reduction

By a gcd argument, we have

**Proposition 3:** Let  $P_1, P_2, Q_1, Q_2$  be polynomials specified in Problem 1. Set

$$\tilde{P}_i(z, w) = w^n P_i\left(\frac{z}{w}\right), \quad \text{and} \quad \tilde{Q}_i(z, w) = w^n Q_i\left(\frac{z}{w}\right),$$

which are homogeneous of degree  $n$  in  $\mathbb{K}[z, w]$ ,  $i = 1, 2$ . Assume  $\frac{A(x)}{B(x)}$  is a solution of (3), where  $A, B \in \mathbb{K}[x]$  with  $\gcd(A, B) = 1$ . Then there exists  $c \in \mathbb{K}$  s.t.

$$\begin{cases} \tilde{P}_1(A(x+1), B(x+1)) = c \cdot \tilde{P}_2(A(x), B(x)), \\ \tilde{Q}_1(A(x+1), B(x+1)) = c \cdot \tilde{Q}_2(A(x), B(x)). \end{cases} \quad (4)$$

By doing coefficient comparison, we can determine **finite** candidates for  $c$  algorithmically. WLOG, we assume that  $c = 1$ .

# Uncoupling

$$\begin{cases} \tilde{P}_1(A(x+1), B(x+1)) = \tilde{P}_2(A(x), B(x)), \\ \tilde{Q}_1(A(x+1), B(x+1)) = \tilde{Q}_2(A(x), B(x)). \end{cases} \quad (5)$$

# Uncoupling

$$\begin{cases} \tilde{P}_1(A(x+1), B(x+1)) = \tilde{P}_2(A(x), B(x)), \\ \tilde{Q}_1(A(x+1), B(x+1)) = \tilde{Q}_2(A(x), B(x)). \end{cases} \quad (5)$$

Applying  $\sigma : x \mapsto x + 1$  to the above equations, we get

$$\begin{cases} \tilde{P}_1(A(x+2), B(x+2)) = \tilde{P}_2(A(x+1), B(x+1)), \\ \tilde{Q}_1(A(x+2), B(x+2)) = \tilde{Q}_2(A(x+1), B(x+1)). \end{cases} \quad (6)$$

Regarding  $A(x+i)$  and  $B(x+i)$  as undeterminates, we have **4 equations** and **6 variables**. It is possible to utilize nonlinear elimination techniques to eliminate **3 variables**, *i.e.*,  $A(x+i)$ 's or  $B(x+i)$ 's from (5) and (6).

# Uncoupling

**Algorithm 1:** Given the difference system (4). Compute **nonzero** autonomous second-order AODEs for  $A(x)$  and  $B(x)$ , respectively, which are consequences of (4).

(1) Let  $I \subseteq \mathbb{K}[w_0, w_1, w_2, z_0, z_1, z_2]$  be the ideal generated by

$$\begin{aligned} &\tilde{P}_1(z_1, w_1) - \tilde{P}_2(z_0, w_0), \quad \tilde{Q}_1(z_1, w_1) - \tilde{Q}_2(z_0, w_0), \\ &\tilde{P}_1(z_2, w_2) - \tilde{P}_2(z_1, w_1), \quad \tilde{Q}_1(z_2, w_2) - \tilde{Q}_2(z_1, w_1). \end{aligned}$$

Using Gröbner bases or resultants, compute nonzero elements  $F_A \in I \cap \mathbb{K}[z_0, z_1, z_2]$  and  $F_B \in I \cap \mathbb{K}[w_0, w_1, w_2]$ .

(2) Return  $F_A(A(x), A(x+1), A(x+2)) = 0$  and  $F_B(B(x), B(x+1), B(x+2)) = 0$ .

# Uncoupling

**Theorem 3 (Vo-Z. 2020)** The elimination ideals  $I \cap \mathbb{K}[z_0, z_1, z_2]$  and  $I \cap \mathbb{K}[w_0, w_1, w_2]$  are nonzero and Algorithm 1 is correct.

## Polynomial solutions

Let  $\frac{A(x)}{B(x)}$  be a solution of the autonomous separable O $\Delta$ E. By Algorithm 1, we can find **nonzero** autonomous second-order AO $\Delta$ Es for  $A(x)$  and  $B(x)$ , respectively.

**Question 2:** Let  $F \in \mathbb{K}[y, z, w]$  be a homogeneous polynomial. Find all polynomial solutions of the AO $\Delta$ E

$$F(y(x), y(x+1), y(x+2)) = 0. \quad (7)$$

## Polynomial solutions

Let  $\frac{A(x)}{B(x)}$  be a solution of the autonomous separable O $\Delta$ E. By Algorithm 1, we can find **nonzero** autonomous second-order AO $\Delta$ Es for  $A(x)$  and  $B(x)$ , respectively.

**Question 2:** Let  $F \in \mathbb{K}[y, z, w]$  be a homogeneous polynomial. Find all polynomial solutions of the AO $\Delta$ E

$$F(y(x), y(x+1), y(x+2)) = 0. \quad (7)$$

**Idea:** Doing coefficient comparison to derive a degree bound.



## Polynomial solutions

Let  $\frac{A(x)}{B(x)}$  be a solution of the autonomous separable O $\Delta$ E. By Algorithm 1, we can find **nonzero** autonomous second-order AO $\Delta$ Es for  $A(x)$  and  $B(x)$ , respectively.

**Question 2:** Let  $F \in \mathbb{K}[y, z, w]$  be a homogeneous polynomial. Find all polynomial solutions of the AO $\Delta$ E

$$F(y(x), y(x+1), y(x+2)) = 0. \quad (7)$$

**Idea:** Doing coefficient comparison to derive a degree bound. Note that (7) is equivalent to

$$\tilde{F}(y(x), \Delta y(x), \Delta^2 y(x)) = 0, \quad (8)$$

where  $\Delta y(x) = y(x+1) - y(x)$  and

$$\tilde{F}(y, z, w) = F(y, y+z, y+2z+w).$$

## Polynomial solutions

For  $\mathbf{i} = (i_1, i_2, i_3) \in \mathbb{N}^3$ , we define  $\|\mathbf{i}\| = i_1 + i_2 + i_3$ . Write

$$\tilde{F} = \sum_{\|\mathbf{i}\|=D} c_{\mathbf{i}} y^{i_1} z^{i_2} w^{i_3},$$

where  $c_{\mathbf{i}} \in \mathbb{K}$ . Set

$$\mathcal{E}(\tilde{F}) = \{\mathbf{i} \in \mathbb{N}^3 \mid c_{\mathbf{i}} \neq 0\},$$

$$m(\tilde{F}) = \min\{i_2 + 2i_3 \mid \mathbf{i} \in \mathcal{E}(\tilde{F})\},$$

$$\mathcal{M}(\tilde{F}) = \{\mathbf{i} \in \mathcal{E}(\tilde{F}) \mid i_2 + 2i_3 = m(\tilde{F})\},$$

$$\mathcal{P}_{\tilde{F}}(t) = \sum_{\mathbf{i} \in \mathcal{M}(\tilde{F})} c_{\mathbf{i}} t^{i_2} [t(t-1)]^{i_3}.$$

We call  $\mathcal{P}_{\tilde{F}}(t)$  the **indicial polynomial** of  $\tilde{F}$  (at infinity).

# Polynomial solutions

**Proposition 4:** Let  $\mathcal{P}_{\tilde{F}}(t)$  be the indicial polynomial of  $\tilde{F}$  at infinity. Then  $\mathcal{P}_{\tilde{F}}(t) \neq 0$ .

**Theorem 4 (Vo-Z. 2020):** Let  $p(x)$  be a nonzero polynomial solution of  $\tilde{F}(y(x), \Delta y(x), \Delta^2 y(x)) = 0$  with degree  $d$ . Then  $\mathcal{P}_{\tilde{F}}(d) = 0$ .

# Algorithms

**Algorithm 2:** Given a separable AO $\Delta$ E  $\frac{P_1(y(x+1))}{Q_1(y(x+1))} = \frac{P_2(y(x))}{Q_2(y(x))}$  with  $\gcd(P_i, Q_i) = 1$  and  $\deg \frac{P_1}{Q_1} = \deg \frac{P_2}{Q_2} \geq 1$ ,  $i = 1, 2$ . Compute a degree bound for its rational solutions.

- (1) Let  $\tilde{P}_j(z, w) = w^n P_j\left(\frac{z}{w}\right)$  and  $\tilde{Q}_j(z, w) = w^n Q_j\left(\frac{z}{w}\right)$ ,  $j = 1, 2$ . Consider

$$\begin{cases} \tilde{P}_1(A(x+1), B(x+1)) = \tilde{P}_2(A(x), B(x)), \\ \tilde{Q}_1(A(x+1), B(x+1)) = \tilde{Q}_2(A(x), B(x)), \end{cases} \quad (9)$$

where  $A, B$  are unknown. Derive the following nonzero AO $\Delta$ Es for  $A(x)$  and  $B(x)$  from (9) by using Algorithm 1:

$$F_A(A(x), A(x+1), A(x+2)) = 0, F_B(B(x), B(x+1), B(x+2)) = 0.$$

# Algorithms

- (2) Determine the indicial polynomials  $\mathcal{P}_{F_A}$  and  $\mathcal{P}_{F_B}$  of  $F_A$  and  $F_B$ , respectively. Let

$$D_A = \{\text{non-negative integer solutions of } \mathcal{P}_{F_A}(t)\},$$

$$D_B = \{\text{non-negative integer solutions of } \mathcal{P}_{F_B}(t)\}.$$

Return  $\max(D_A \cup D_B)$ .

# Algorithms

**Algorithm 3:** Given an irreducible autonomous first-order AODE  $F(y(x), y(x+1)) = 0$ . Compute a non-constant rational solution or return “NULL”.

- (1) If  $\deg_y(F) \neq \deg_z(F)$ , then output “NULL”. Otherwise, go to step 2.
- (2) Compute the genus  $g$  of the corresponding curve  $\mathcal{C}_F$  defined by  $F(y, z) = 0$ . If  $g \neq 0$ , then output “NULL”. Otherwise, go to step 3.
- (3) Using Vo-Grasegger-Winkler’s algorithm, determine an optimal parametrization for  $\mathcal{C}_F$ , say  $\mathcal{P}(t) = (p_1(t), p_2(t))$ .

# Algorithms

- (4) Apply Algorithm 2 to compute a degree bound  $N$  for rational solutions of the separable AODE  $p_1(y(x+1)) = p_2(y(x))$ .
- (5) Set  $M = N \cdot \deg p_1$ . Use Feng-Gao-Huang's algorithm to determine a non-constant rational solution of  $F(y) = 0$  whose degree is at most  $M$ . Return the rational solution if there is any. Otherwise, return "NULL".

## Example

Consider the first-order autonomous AOΔE:

$$F = (12y(x) + 49)y(x + 1)^2 - (12y^2 + 62y + 56)y(x + 1) + y(x)^2 + 8y(x) + 16 = 0. \quad (10)$$

It is clear that  $\deg_y(F) = \deg_z(F) = 2$ . The corresponding algebraic curve is of genus zero and it has an optimal parametrization

$$\mathcal{P}(t) = (p_1(t), p_2(t)) = \left( \frac{9t^2 - 12t + 4}{12t}, \frac{9t^2 + 36t + 4}{12(t + 4)} \right).$$

Using the above parametrization, we derive the following associated separable AOΔE of (10):

$$\frac{9y(x + 1)^2 - 12y(x + 1) + 4}{y(x + 1)} = \frac{9y(x)^2 + 36y(x) + 4}{y(x) + 4}. \quad (11)$$



## Example

Using Algorithm 2, we see the degree bound for rational solutions of (11) is 2. Thus, the degrees of rational solutions of  $F(y) = 0$  are bounded by 4. Applying Feng-Gao-Huang's algorithm, we determine a rational solution, say

$$y(x) = \frac{(1 - 4x + 2x^2)^2}{2x(1 - 3x + 2x^2)}.$$

# Conclusion

- ▶ An algebraic geometric approach for studying rational solutions of first-order AODEs.
- ▶ A degree bound for rational solutions of autonomous first-order AODEs, and thus derive a complete algorithm for computing corresponding rational solutions.

# Conclusion

- ▶ An algebraic geometric approach for studying rational solutions of first-order AODEs.
- ▶ A degree bound for rational solutions of autonomous first-order AODEs, and thus derive a complete algorithm for computing corresponding rational solutions.

Thanks!